

ARTICLE

Every graph occurs as an induced subgraph of some hypohamiltonian graph

Carol T. Zamfirescu¹  | Tudor I. Zamfirescu^{2,3,4}

¹Department of Applied Mathematics, Computer Science and Statistics, Ghent University, Krijgslaan 281 - S9, 9000 Ghent, Belgium

²Fachbereich Mathematik, Universität Dortmund, 44221 Dortmund, Germany

³“Simion Stoilow” Institute of Mathematics, Roumanian Academy, Bucharest, Roumania

⁴College of Mathematics and Information Science, Hebei Normal University, 050024 Shijiazhuang, P.R. China

Correspondence

Carol T. Zamfirescu, Department of Applied Mathematics, Computer Science and Statistics, Ghent University, Krijgslaan 281 - S9, 9000 Ghent, Belgium.
Email: czamfirescu@gmail.com

Funding information

Fonds Wetenschappelijk Onderzoek

Abstract

We prove the titular statement. This settles a problem of Chvátal from 1973 and encompasses earlier results of Thomassen, who showed it for K_3 , and Collier and Schmeichel, who proved it for bipartite graphs. We also show that for every outerplanar graph there exists a planar hypohamiltonian graph containing it as an induced subgraph.

KEYWORDS

hypohamiltonian, induced subgraph

MSC 2010.

05C10, 05C45, 05C60

1 | INTRODUCTION

Consider a non-hamiltonian graph G . We call G *hypohamiltonian* if for every vertex v in G , the graph $G - v$ is hamiltonian. In similar spirit, G is said to be *almost hypohamiltonian* if there exists a vertex w in G , which we will call *exceptional*, such that $G - w$ is non-hamiltonian, and for any vertex $v \neq w$ in G , the graph $G - v$ is hamiltonian. For an overview of results on hypohamiltonicity till 1993, see the survey by Holton and Sheehan [7]. For newer material that also includes work on the recently introduced almost hypohamiltonian graphs, we refer the reader to [4,5,8,15] and the references found therein.

In 1973, Chvátal [2] asked whether every graph may occur as the induced subgraph of some hypohamiltonian graph. As Thomassen writes in [10], an important partial answer was provided by Collier and Schmeichel [3] who proved that every bipartite graph is an induced subgraph of some hypohamiltonian graph. In [11], Thomassen constructs an infinite family of planar cubic hypohamiltonian graphs and shows that certain edges can be added to these graphs such that the resulting graphs are hypohamiltonian, as well. He uses this to give a simple proof of the aforementioned result of Collier and Schmeichel.

Earlier, Thomassen [9] had proven that hypohamiltonian graphs of girth 3 and 4 exist, i.e. that K_3 and C_4 can occur as induced subgraphs of hypohamiltonian graphs—see also [3]. This refuted the conjecture of Herz et al. [6] that hypohamiltonian graphs have girth at least 5. Thomassen emphasizes in [10] that even for K_4 the answer to Chvátal's problem is unknown.

In this note, we prove that *any* graph can appear as an induced subgraph of some hypohamiltonian graph.

2 | AUXILIARY RESULTS

Consider a planar almost hypohamiltonian graph with a cubic exceptional vertex, for example the graph F of order 36 (discovered by Goedgebeur and the first author [5], and independently by Wiener [13, 14]), with exceptional vertex v . Note that F is the smallest known planar graph fit for the construction to come—however, there might be smaller graphs usable that have not been found yet. In fact, there is a smaller planar almost hypohamiltonian graph known (found by Wiener [14] and of order 31), but it does not contain a cubic exceptional vertex, which is needed for the method to work.




FIGURE 1 The copy F'' of F . The exceptional vertex is marked v''

Take two disjoint copies F', F'' of F , with $v' \in V(F')$ and $v'' \in V(F'')$ corresponding to v . Put $N(v') = \{o, b, c\}$ and $N(v'') = \{o', d, a\}$. For an illustration of F'' , see Figure 1. Take $F' - v'$, $F'' - v''$, identify o with o' , and add the edges ab and cd . The neighbors a, b, c, d are chosen such that we obtain the graph H depicted in Figure 2. (The “half-edges” shown in Figures 2 and 3 end in vertices outside H .) Already Thomassen used such a construction in [9]. In what follows, we see $F' - v'$ and $F'' - v''$ as subgraphs of H . In particular, a, b, c, d denote vertices in H , as well.

Suppose an arbitrary but fixed graph W has a hamiltonian cycle Λ , and let W include the graph H such that among the vertices of H only a, b, c, d are connected by edges with vertices in $W - H$.

Lemma 1. *Either $H \cap \Lambda$ is (i) the union of two disjoint paths, one from a to b and the other from c to d , or (ii) a path from a to c , or (iii) a path from b to d .*

Proof. There is no hamiltonian path \mathfrak{p} in H between a and b , since \mathfrak{p} would have to use the edge dc , yielding a hamiltonian path between a and d in $F'' - v''$ or a hamiltonian path between c and b in $F' - v'$ (depending on when \mathfrak{p} picks up o). Adding the path $av''d$ or $bv'c$, respectively, we get a hamiltonian cycle in F and a contradiction is obtained. Now assume there is a hamiltonian path \mathfrak{q} in




FIGURE 2 We depict diagrammatically the graph H shown above with the rectangle shown below



FIGURE 3 There are two essentially different ways to traverse the rectangle $abcd$: in the first situation $(ab \cup cd)$ we traverse disconnectedly, while in the latter two $(ac \text{ or } bd)$ we traverse diagonally

H between a and d . Then there exists a hamiltonian path between o and b in F' or a hamiltonian path between o and c in F' . As above, we are led to a contradiction since F is non-hamiltonian.

If $H \cap \Lambda$ is the union of two disjoint paths, one from b to c , the other from d to a , then one of these paths must contain o . Assume w.l.o.g. the former to be that path. Considering it in F' and adding to it the path $bu'c$, we obtain a hamiltonian cycle in F' , a contradiction since F' is almost hypohamiltonian. ■

In case (i) we say that H is *disconnectedly traversed*, while if case (ii) or (iii) occurs H is called *diagonally traversed*. In cases (ii) and (iii) of Lemma 1, when H is diagonally traversed, we say more precisely that H is $a - c$ and $b - d$ traversed, respectively.

Consider the graph G^* of Figure 4. There, A, B, C, \dots, Q are graphs isomorphic to H . The length of the cycle $\Gamma = wy y^* \dots y'$ equals the number of copies of H used to construct G^* , i.e. $|\{A, B, C, \dots, Q\}| = |V(\Gamma)|$. Γ is included in an arbitrary hamiltonian graph Z with the hamiltonian cycle Γ .

Lemma 2. *The graph G^* is not hamiltonian.*

Proof. Suppose G^* has a hamiltonian cycle Λ . Obviously, not all copies of H are disconnectedly traversed. Suppose A is $a - c$ traversed. Then Λ quickly visits y , as it must continue with the path $cuxy$. If B is also diagonally traversed, then analogously Λ quickly visits y , and Λ is not hamiltonian. Hence, by Lemma 1, B is disconnectedly traversed. Thus $xyz \subset \Lambda$.




FIGURE 4 The graph G^*

Necessarily, Q is $b-d$ traversed. So, analogously, $x'y'z' \subset \Lambda$. Continuing this reasoning, we see that no edge of Z is in Λ . Indeed, all other copies of H are disconnectedly traversed, so $x^*y^*z^* \subset \Lambda$ etc. Thus $w \notin \Lambda$, and a contradiction is obtained. ■

3 | MAIN RESULTS

A graph is *outerplanar* if it possesses a planar embedding in which every vertex belongs to the unbounded face. Note that a graph is outerplanar if and only if it does not contain a graph homeomorphic to K_4 or $K_{2,3}$, see [1]. We now present our main theorem.

Theorem 1. *Every graph is contained in some hypohamiltonian graph as an induced subgraph.*

Proof. Let G be an arbitrary graph. In the remainder of this proof, all notation refers to notions introduced in Section 2. Choose Z to be Γ to which G is added in such a way that the finite sequence of its vertices is placed at every second vertex of Γ . (So, the length of Γ is $2|V(G)|$.) By Lemma 2, G^* is not hamiltonian. It remains to provide a hamiltonian cycle in $G^* - s$, for every vertex s in G^* .

A hamiltonian cycle of $G^* - w$ is shown in Figure 4. By changing uxy into uwv we get a hamiltonian cycle in $G^* - x$.

Due to the symmetries, it remains to show that $G^* - s$ is hamiltonian for any $s \in V(F')$. Consider $s \in V(F')$. There is a hamiltonian path in $F' - s$ joining b to c , or c to o , or o to b . In the second case (c to o), we change the route of Λ inside the subgraph spanned by $V(A) \cup \{x, y, u, w\}$ as shown in Figure 5 (a), and in the third case (o to b), we change the route as depicted in Figure 5 (b). In the first case, if s is the central vertex o of A , we obtain a hamiltonian cycle of $G^* - o$ as pictured in Figure 6. For other positions of $s \in V(F')$, see Figure 7. ■

From the above proof, we immediately obtain the following.




FIGURE 5 The route of Λ inside the subgraph spanned by $V(A) \cup \{x, y, u, w\}$




FIGURE 6 A hamiltonian cycle of $G^* - o$




FIGURE 7 A hamiltonian cycle of $G^* - s$

Theorem 2. *If G is an outerplanar graph, then there exists a planar hypohamiltonian graph containing G as an induced subgraph.*

Theorem 2 cannot be extended to include *all* planar graphs due to an elegant argument of Thomassen, who proves in [11] that by Whitney's Theorem [12]—which states that planar triangulations without separating triangles are hamiltonian—, a planar triangulation cannot be an induced subgraph of any pla-

nar hypohamiltonian graph. However, by Theorem 1, any planar triangulation is very well an induced subgraph of some (necessarily nonplanar) hypohamiltonian graph.

Thomassen's results from [11] can be used to describe certain bipartite planar graphs that can be subgraphs of planar hypohamiltonian graphs. He goes on to write: "Maybe this is the case for every bipartite planar graph." This question remains unresolved, as for instance $K_{2,3}$ is planar yet neither a planar triangulation, nor among Thomassen's aforementioned planar bipartite graphs, nor an outerplanar graph.

Theorem 3.

- (i) *There exists a hypohamiltonian graph of order $20n$ containing K_n as an induced subgraph. In particular, there exists a hypohamiltonian graph of order 80 containing K_4 .*
- (ii) *There exists a planar hypohamiltonian graph of girth 3 and order 216.*
- (iii) *For an outerplanar graph G of order n there exists a planar hypohamiltonian graph of order $144n$ containing G as an induced subgraph.*

Proof. For (i), we may modify the construction from the proof of Theorem 1 by using every vertex of Γ (since G is in this case a complete graph). Thus, here the length of Γ is $|V(K_n)| = n$. Furthermore, we use the Petersen graph instead of the graph from Figure 1. It is now easy to verify that the order of G^* is indeed $20n$.

The proof of (ii) is the same as the proof of (i), but we replace Petersen's graph by the plane almost hypohamiltonian graph shown in Figure 1. Here, $n = 3$.

In (iii), the length of Γ is $2n$. We use $2n$ copies of the graph from Figure 2, which is of order 69. We obtain a graph of order $((69 + 4) \cdot 2n) - 2n = 144n$. ■

Part (ii) improves a bound given in [4, Corollary 3.4]. We end this note with the following.

Problem. Characterize those planar graphs that occur as induced subgraphs of planar hypohamiltonian graphs.

ACKNOWLEDGMENT

The research of Carol T. Zamfirescu is supported by a Postdoctoral Fellowship of the Research Foundation Flanders (FWO).

ORCID

Carol T. Zamfirescu  <http://orcid.org/0000-0002-9673-410X>

REFERENCES

- [1] G. Chartrand and F. Harary, *Planar permutation graphs*, Ann. Inst. Henri Poincaré, Nouv. Sér. Sect. B **3** (1967), 433–438.
- [2] V. Chvátal, *Flip-flops in hypohamiltonian graphs*, Canad. Math. Bull. **16** (1973), 33–41.
- [3] J. B. Collier and E. F. Schmeichel, *New flip-flop constructions for hypohamiltonian graphs*, Discrete Math. **18** (1977), 265–271.
- [4] J. Goedgebeur and C. T. Zamfirescu, *Improved bounds for hypohamiltonian graphs*, Ars Math. Contemp. **13** (2017), 235–257.
- [5] J. Goedgebeur and C. T. Zamfirescu, *On almost hypohamiltonian graphs*. Submitted. See Section 3 of arXiv:1606.06577 [math.CO].

- [6] J. C. Herz, J. J. Duby, and F. Vigué, Recherche systématique des graphes hypohamiltoniens, in: *Theory of Graphs: International Symposium*, Rome, 1966, pp. 153–159, Gordon and Breach, New York, and Dunod, Paris, 1967.
- [7] D. A. Holton and J. Sheehan, *The Petersen graph, Chapter 7: Hypohamiltonian graphs*, Cambridge University Press, New York, 1993.
- [8] M. Jooyandeh et al., *Planar hypohamiltonian graphs on 40 vertices*, *J. Graph Theory* **84** (2017), 121–133.
- [9] C. Thomassen, *On hypohamiltonian graphs*, *Discrete Math.* **10** (1974), 383–390.
- [10] C. Thomassen, *Hypohamiltonian graphs and digraphs. Theory and Applications of Graphs*, *Lecture Notes in Mathematics*, **642**, Springer, Berlin (1978), pp. 557–571.
- [11] C. Thomassen, *Planar cubic hypohamiltonian and hypotractable graphs*, *J. Combin. Theory Ser. B* **30** (1981), 36–44.
- [12] H. Whitney, *A theorem on graphs*, *Ann. Math.* **32** (1931), 378–390.
- [13] G. Wiener, *On constructions of hypotractable graphs*, *Electron. Notes Discrete Math.* **54** (2016), 127–132.
- [14] G. Wiener, *New constructions of hypohamiltonian and hypotractable graphs*, To appear in *J. Graph Theory*, <https://doi.org/10.1002/jgt.22173>
- [15] C. T. Zamfirescu, *Hypohamiltonian and almost hypohamiltonian graphs*, *J. Graph Theory* **79** (2015), 63–81.

How to cite this article: Zamfirescu CT, Zamfirescu TI. Every graph occurs as an induced subgraph of some hypohamiltonian graph. *J Graph Theory*. 2017;00:1–7. <https://doi.org/10.1002/jgt.22228>